

THE MILNOR INVARIANTS OF CLOVER LINKS

KODAI WADA AND AKIRA YASUHARA

ABSTRACT. J.P. Levine introduced a clover link to investigate the indeterminacy of the Milnor invariants of a link. It is shown that for a clover link, the Milnor numbers of length at most $2k + 1$ are well-defined if those of length at most k vanish, and that the Milnor numbers of length at least $2k + 2$ are not well-defined if those of length $k + 1$ survive. For a clover link c with the Milnor numbers of length at most k vanishing, we show that the Milnor number $\mu_c(I)$ for a sequence I is well-defined up to the greatest common divisor of $\mu_c(J)$'s, where J is a subsequence of I obtained by removing at least $k + 1$ indices. Moreover, if I is a non-repeated sequence with length $2k + 2$, the possible range of $\mu_c(I)$ is given explicitly. As an application, we give an edge-homotopy classification of 4-clover links.

1. INTRODUCTION

The *Milnor invariant* introduced by J. Milnor [7], [8]. For an oriented ordered n -component link L in the 3-sphere S^3 with peripheral information, the *Milnor number* $\mu_L(I)$, which is an integer, is specified by a finite sequence I in $\{1, 2, \dots, n\}$. The Milnor $\bar{\mu}$ -invariant $\bar{\mu}_L(I)$ is the residue class of $\mu_L(I)$ modulo the greatest common divisor of $\mu_L(J)$'s, where J is obtained from proper subsequence of I by permuting cyclicly. The length of the sequence I is called the *length* of $\bar{\mu}_L(I)$ and denoted by $|I|$. His original definition of the Milnor invariant eliminates the indeterminacy of the possible variations of the Milnor numbers caused by different choices of peripheral elements.

In [6], J. P. Levine examined the Milnor invariants from the point of view of *based* links, in order to understand the indeterminacy. A based link is a link for which some peripheral information is specified, i.e., meridians (the weakest) or both meridians and longitudes (the strongest). It is known that these invariants are completely well-defined for the strongest form of basing (disk links [5] or string links [2]). As basing only slightly stronger than the specification of meridians, he introduced an *n-clover link* which is an embedded graph consisting of n loops, each loop connected to a vertex by an edge in S^3 . The Milnor number of a clover link is defined up to flatly isotopy and it is shown that those of length $\leq k$ vanish implies those of length $\leq 2k + 1$ are completely non-indeterminate [6].

The first author [10] redefined the Milnor number of a clover link up to ambient isotopy as follows. Given an n -clover link c , we construct an n -component *bottom tangle* $\gamma(F_c)$ by using a *disk/band surface* F_c of c . In [6], Levine defined the Milnor number of a bottom tangle. Therefore we define the *Milnor number* μ_c of an *n-clover link* c to be the Milnor number $\mu_{\gamma(F_c)}$. (In [6], a bottom tangle is called a string link. The name ‘bottom tangle’ follows K. Habiro [3].) In [10], it is shown that the same result as Levine [6] holds while there are infinitely many choices of $\gamma(F_c)$ for c .

Date: July 7, 2015.

The second author is partially supported by a JSPS Grant-in-Aid for Scientific Research (C) (#26400081).

Unfortunately, the Milnor numbers of length at least $2k+2$ are not well-defined if those of length $k+1$ survive. In this paper, we show that the Milnor number $\mu_c(I)$ of a sequence I modulo $\delta_c^k(I)$ is well-defined if the Milnor numbers of length $\leq k$ vanish, where $\delta_c^k(I)$ is the greatest common divisor of $\mu_c(J)$'s, where J is range to over all subsequences of I obtained by removing at least $k+1$ indices. In fact, we have the following theorem.

Theorem 1.1. *Let c be an n -clover link and l_c a link which is the disjoint union of loops of c . If the Milnor numbers of l_c for sequences with length $\leq k$ vanish, then the residue class of $\mu_c(I)$ modulo $\delta_c^k(I)$ is an invariant for any sequence I .*

Remark 1.2. *In contrast to the $\bar{\mu}$ -invariant for a link in S^3 , we do not need to take cyclic permutation for getting $\delta_c^k(I)$.*

It is an important property that for non-repeated sequences, the Milnor invariants of links are link-homotopy invariants [7]. The first author showed that the Milnor numbers for any non-repeated sequence with length ≤ 3 give an *edge-homotopy* classification of 3-clover links [10], where edge-homotopy [9] is an equivalence relation, which is a generalization of link-homotopy, generated by crossing changes on the same spatial edge. We also discuss giving an edge-homotopy classification of 4-clover links. The Milnor numbers for non-repeated sequences with length 4 could be useful to have an edge-homotopy classification of 4-clover links. But they are not well-defined in general. Hence we consider the set of all Milnor numbers of length 4 for all disk/band surfaces of c . More generally, we define the following set;

$$H_c(2k+2, j) = \left\{ \sum_{S \in \mathcal{S}_j^{2k+1}} \mu_{\gamma(F_c)}(Sj) X_S \mid F_c : \text{a disk/band surface of } c \right\}$$

for each integer j ($1 \leq j \leq n$), where $\mathcal{S}_j^{2k+1}$ is the set of length- $(2k+1)$ non-repeated sequences without containing j and for a sequence $S = i_1 i_2 \dots i_{2k+1}$, $X_S = X_{i_1} X_{i_2} \dots X_{i_{2k+1}}$ is a monomial in non-commutative variables $X_1, \dots, X_n$. Since $H_c(2k+2, j)$ consists of the Milnor numbers $\mu_{\gamma(F_c)}(Sj)$ ($S \in \mathcal{S}_j^{2k+1}$) for all disk/band surfaces of c , it is an invariant of c . While it seems too big to handle $H_c(2k+2, j)$, we have the following theorem.

Theorem 1.3. *Let c be an n -clover link and F_c a disk/band surface of c . If the Milnor numbers of l_c for non-repeated sequences with length $\leq k$ vanish, then we have the following:*

$$\begin{aligned} H_c(2k+2, j) = & \left\{ \sum_{\substack{|J|=|I|=k \\ JI \in \mathcal{S}_j^{2k+1}}} \mu_{\gamma(F_c)}(Jj) \mu_{\gamma(F_c)}(II) \left(\sum_{i_s \in \{J\}} m_{li_s} X_{J_{<s}} (X_{i_s II} - X_{i_s I I} - X_{II i_s} + X_{II i_s}) X_{J_{s<}} \right. \right. \\ & \left. \left. + m_{lj} (X_{IIJ} - X_{I I J} - X_{J II} + X_{J I I}) \right) + \sum_{S \in \mathcal{S}_j^{2k+1}} \mu_{\gamma(F_c)}(Sj) X_S \mid m_{pq} = m_{qp} \in \mathbb{Z} \right\}, \end{aligned}$$

where for a sequence $J = i_1 \dots i_m$, $\{J\} = \{i_1, \dots, i_m\}$ and $J_{<s}$ (resp. $J_{s<}$) is a subsequence $i_1 \dots i_{s-1}$ (resp. $i_{s+1} \dots i_m$) of J for $1 \leq s \leq m$, and $X_{J_{<1}}$ and $X_{J_{m<}}$ are defined to be 1.

This theorem implies that the set $H_c(2k+2, j)$ is obtained from the Milnor numbers of $\gamma(F_c)$ for any single disk/band surface F_c of c , that is, $H_c(2k+2, j)$ is specified explicitly.

By the following corollary we have that $H_c(2k+2, j)$ is not only an invariant of c but also an edge-homotopy invariant.

Corollary 1.4. *For a clover link c , if the Milnor numbers of l_c for non-repeated sequences with length $\leq k$ vanish, then $H_c(2k+2, j)$ is an edge-homotopy invariant of c .*

It is the definition that the Milnor numbers of length 1 are zero. If $k = 1$, then the theorem above holds without the condition. The following example follows directly from Theorem 1.3.

Example 1.5. *For a 4-clover link c and a disk/band surface F_c of c , we have*

$$\begin{aligned} H_c(4, 4) = & \left\{ \begin{aligned} & \left(\begin{array}{l} \mu_{\gamma(F_c)}(14)\mu_{\gamma(F_c)}(23)(m_{13} - m_{34} - m_{12} + m_{24}) \\ + \mu_{\gamma(F_c)}(12)\mu_{\gamma(F_c)}(34)(m_{24} - m_{23} - m_{14} + m_{13}) \end{array} \right) X_{123} \\ & + \left(\begin{array}{l} \mu_{\gamma(F_c)}(14)\mu_{\gamma(F_c)}(23)(m_{34} - m_{13} - m_{24} + m_{12}) \\ + \mu_{\gamma(F_c)}(13)\mu_{\gamma(F_c)}(24)(m_{34} - m_{23} - m_{14} + m_{12}) \end{array} \right) X_{132} \\ & + \left(\begin{array}{l} \mu_{\gamma(F_c)}(13)\mu_{\gamma(F_c)}(24)(m_{23} - m_{34} - m_{12} + m_{14}) \\ + \mu_{\gamma(F_c)}(12)\mu_{\gamma(F_c)}(34)(m_{23} - m_{24} - m_{13} + m_{14}) \end{array} \right) X_{213} \\ & + \left(\begin{array}{l} \mu_{\gamma(F_c)}(14)\mu_{\gamma(F_c)}(23)(m_{34} - m_{13} - m_{24} + m_{12}) \\ + \mu_{\gamma(F_c)}(13)\mu_{\gamma(F_c)}(24)(m_{34} - m_{23} - m_{14} + m_{12}) \end{array} \right) X_{231} \\ & + \left(\begin{array}{l} \mu_{\gamma(F_c)}(13)\mu_{\gamma(F_c)}(24)(m_{23} - m_{34} - m_{12} + m_{14}) \\ + \mu_{\gamma(F_c)}(12)\mu_{\gamma(F_c)}(34)(m_{23} - m_{24} - m_{13} + m_{14}) \end{array} \right) X_{312} \\ & + \left(\begin{array}{l} \mu_{\gamma(F_c)}(14)\mu_{\gamma(F_c)}(23)(m_{13} - m_{34} - m_{12} + m_{24}) \\ + \mu_{\gamma(F_c)}(12)\mu_{\gamma(F_c)}(34)(m_{24} - m_{23} - m_{14} + m_{13}) \end{array} \right) X_{321} \\ & + \sum_{S \in \mathcal{S}_4^3} \mu_{\gamma(F_c)}(S4)X_S \mid m_{pq} \in \mathbb{Z} \end{aligned} \right\}. \end{aligned}$$

This together with the theorem below gives us an edge-homotopy classification of 4-clover links, see Remark 1.7.

Theorem 1.6. *Let c and c' be 4-clover links. They are edge-homotopic if and only if $H_c(4, 4) \cap H_{c'}(4, 4) \neq \emptyset$ and $\mu_c(I) = \mu_{c'}(I)$ for any non-repeated sequence I with $|I| \leq 3$.*


Remark 1.7. *By Example 1.5, we are able to determine whether $H_c(4, 4) \cap H_{c'}(4, 4)$ is empty or not. Hence by combining Example 1.5 and Theorem 1.6, we obtain an edge-homotopy classification of 4-clover links.*

2. THE MILNOR NUMBERS OF CLOVER LINKS

In this section we define the Milnor numbers for clover links.

2.1. A construction of bottom tangles. An n -component *tangle* is a properly embedded disjoint union of n arcs in the 3-cube $[0, 1]^3$. An n -component *bottom tangle* $\gamma = \gamma_1 \cup \gamma_2 \cup \dots \cup \gamma_n$ defined by Levine [6] is a tangle with $\partial\gamma_i = \{(\frac{2i-1}{2n+1}, \frac{1}{2}, 0), (\frac{2i}{2n+1}, \frac{1}{2}, 0)\} \subset \partial[0, 1]^3$ for each i ($= 1, 2, \dots, n$).

A *spatial graph* is an embedded graph in S^3 . Let C_n be a graph consisting of n oriented loops $e_1, e_2, \dots, e_n$, each loop e_i connected to a vertex v by an edge f_i ($i = 1, 2, \dots, n$), see Figure 2.1. An n -clover link in S^3 is a spatial graph of C_n [6]. The each part of a clover link corresponding to e_i , f_i and v of C_n are called the *leaf*, *stem* and *root*, denoted by the same notations respectively.

FIGURE 2.1. the graph C_n

L. Kauffman, J. Simon, K. Wolcott and P. Zhao [4] defined disk/band surfaces of spatial graphs. For a spatial graph Γ , a *disk/band surface* F_Γ of Γ is a compact, oriented surface in S^3 such that Γ is a deformation retract of F_Γ contained in the interior of F_Γ . Note that any disk/band surface of a spatial graph is ambient isotopic to a surface constructed by putting a disk at each vertex of the spatial graph, connecting the disks with bands along the spatial edges. We remark that for a spatial graph, there are infinitely many disk/band surfaces up to ambient isotopy.

Given an n -clover link, we construct an n -component bottom tangle using a disk/band surface of the clover link as follows:

- (1) For an n -clover link c , let F_c be a disk/band surface of c and let D be a disk which contains the root. From now on, we may assume that the intersection

$D \cap \bigcup_{i=1}^n f_i$ and orientations of the disks are as illustrated in Figure 2.2.




FIGURE 2.2.

- (2) Let $N(D)$ be the regular neighborhood of D and $N^\circ(D)$ the interior of $N(D)$. Since $S^3 \setminus N^\circ(D)$ is homeomorphic to the 3-ball, $F_c \setminus N^\circ(D)$ can be seen as a disjoint union of surfaces in the 3-ball. Hence $\partial F_c \setminus N(D)$ is a disjoint union of n -arcs and n -circles $\bigcup_{i=1}^n S_i^1$ in the 3-ball.
- (3) Since the 3-ball is homeomorphic to $[0, 1]^3$, we obtain an oriented ordered n -component bottom tangle $\gamma(F_c)$ from $(\partial F_c \setminus N^\circ(D)) \setminus \bigcup_{i=1}^n S_i^1$ as illustrated in (3) and (4) of Figure 2.3. We call $\gamma(F_c)$ an n -component bottom tangle obtained from F_c .




FIGURE 2.3. A method for obtaining a bottom tangle from a disk-band surface of a clover link

While there are infinitely many disk/band surfaces of a clover link which satisfy the condition (1) above, they are related by certain local moves as follows.

Lemma 2.1. [10, Proposition 2.5] *For an n -clover link c , any two disk/band surfaces F_c and F'_c are transformed into each other by adding full-twists to bands (Figure 2.4 (a)) and a single move illustrated in Figure 2.4 (b).*




FIGURE 2.4. Two local moves of disk/band surfaces

2.2. Milnor invariants. Let us briefly recall from [6] the definition of the Milnor number of a bottom tangle. Let $\gamma = \gamma_1 \cup \gamma_2 \cup \cdots \cup \gamma_n$ be an oriented ordered n -component bottom tangle in $[0, 1]^3$ with $\partial\gamma_i = \{(\frac{2i-1}{2n+1}, \frac{1}{2}, 0), (\frac{2i}{2n+1}, \frac{1}{2}, 0)\} \subset \partial[0, 1]^3$ for each $i (= 1, 2, \dots, n)$. Let G be the fundamental group of $[0, 1]^3 \setminus \gamma$ with a base point $p = (\frac{1}{2}, 0, 0)$ and G_q the q th lower central subgroup of G . Let α_i and λ_i be the i th meridian and i th longitude of γ respectively as illustrated in Figures 2.5. We assume that λ_i is trivial in G/G_2 . Since the quotient group G/G_q is generated by $\alpha_1, \alpha_2, \dots, \alpha_n$ [1], γ is represented by $\alpha_1, \alpha_2, \dots, \alpha_n$ modulo G_q .




FIGURE 2.5. meridians and longitudes

We consider the *Magnus expansion* of λ_j . The Magnus expansion is a homomorphism (denoted E) from a free group $\langle \alpha_1, \alpha_2, \dots, \alpha_n \rangle$ to the formal power series ring in non-commutative variables $X_1, X_2, \dots, X_n$ with integer coefficients defined as follows. $E(\alpha_i) = 1 + X_i$, $E(\alpha_i^{-1}) = 1 - X_i + X_i^2 - X_i^3 + \cdots$ ($i = 1, 2, \dots, n$).

For a sequence $I = i_1 i_2 \dots i_{k-1} j$ ($i_m \in \{1, 2, \dots, n\}, k \leq q$), we define the *Milnor number* $\mu_\gamma(I)$ to be the coefficient of $X_{i_1 i_2 \dots i_{k-1} j}$ in $E(\lambda_j)$ (we define $\mu_\gamma(j) = 0$), which is an invariant [6]. (In [6], the set of λ_i 's, without taking the Magnus expansion, is called the *Milnor's $\bar{\mu}$ -invariant*.) For a bottom tangle $\gamma = \gamma_1 \cup \gamma_2 \cup \cdots \cup \gamma_n$, an oriented link $L(\gamma) = L_1 \cup L_2 \cup \cdots \cup L_n$ in S^3 can be defined by $L_i = \gamma_i \cup a_i$, where a_i is a line segment connecting $(\frac{2i-1}{2n+1}, \frac{1}{2}, 0)$ and $(\frac{2i}{2n+1}, \frac{1}{2}, 0)$, see Figure 2.6. We call $L(\gamma)$ the *closure* of γ . On the other hand, for any oriented link




FIGURE 2.6.

L in S^3 , there is a bottom tangle γ_L such that the closure of γ_L is equal to L . So we define the Milnor number of L to be the Milnor number of γ_L . Let $\Delta_L(I)$ be the greatest common divisor of $\mu_L(J)$'s, where J is obtained from proper subsequence of I by permuting cyclicly. The *Milnor invariant* $\bar{\mu}_L(I)$ is the residue class of $\mu_L(I)$ modulo $\Delta_L(I)$. We note that for a sequence I , if we have $\Delta_L(I) = 0$, then the Milnor invariant $\bar{\mu}_L(I)$ is equal to the Milnor number $\mu_{\gamma_L}(I)$. Now we define the Milnor number of a clover link.

Definition 2.2. Let c be an n -clover link and F_c a disk/band surface of c . Let $\gamma(F_c)$ be the n -component bottom tangle obtained from F_c . For a sequence I , the Milnor number $\mu_c(I)$ of c is defined to be the Milnor number $\mu_{\gamma(F_c)}(I)$.

Remark 2.3. While $\mu_c(I)$ depends on a choice of F_c , the first author [10] proved the following result: Let l_c be a link which is the disjoint union of leaves of c . If the Milnor numbers of l_c for sequences with length $\leq k$ vanish, then $\mu_c(I)$ is well-defined for any sequence I with $|I| \leq 2k + 1$.

3. PROOF OF THEOREM 1.1

In this section we will give a proof of Theorem 1.1.

An n -component tangle $u = u_1 \cup u_2 \cup \dots \cup u_n$ is an n -component *string link* if for each i ($= 1, 2, \dots, n$), the boundary $\partial u_i = \{(\frac{2i-1}{2n+1}, \frac{1}{2}, 0), (\frac{2i-1}{2n+1}, \frac{1}{2}, 1)\} \subset \partial[0, 1]^3$. In particular, u is *trivial* if for each i ($= 1, 2, \dots, n$), $u_i = \{(\frac{2i-1}{2n+1}, \frac{1}{2})\} \times [0, 1]$ in $[0, 1]^3$.

Here we introduce a *SL-move* [10] given by a string link u which is a transformation of an n -component bottom tangle $\gamma = \gamma_1 \cup \gamma_2 \cup \dots \cup \gamma_n$ with $\partial \gamma_i = \{(\frac{2i-1}{2n+1}, \frac{1}{2}, 0), (\frac{2i}{2n+1}, \frac{1}{2}, 0)\} \subset \partial[0, 1]^3$.

- (1) Let $u = u_1 \cup u_2 \cup \dots \cup u_n$ be an oriented ordered n -component string link in $[0, 1]^3$. For each i ($= 1, 2, \dots, n$), we consider an arc u'_i which is parallel to the i th component u_i of u with opposite orientation and $\partial u'_i = \{(\frac{2i}{2n+1}, \frac{1}{2}, 0), (\frac{2i-1}{2n+1}, \frac{1}{2}, 1)\} \subset \partial[0, 1]^3$, see Figure 3.7.




FIGURE 3.7.

- (2) Let $\gamma' = \gamma'_1 \cup \gamma'_2 \cup \dots \cup \gamma'_n$ be an n -component bottom tangle in $[0, 1]^3$ defined by

$$\gamma'_i = h_0(u_i \cup u'_i) \cup h_1(\gamma_i)$$

for $i = 1, 2, \dots, n$, where $h_0, h_1 : ([0, 1] \times [0, 1]) \times [0, 1] \rightarrow ([0, 1] \times [0, 1]) \times [0, 1]$ are embeddings defined by

$$h_0(x, t) = (x, \frac{1}{2}t) \text{ and } h_1(x, t) = (x, \frac{1}{2} + \frac{1}{2}t)$$

for $x \in ([0, 1] \times [0, 1])$ and $t \in [0, 1]$.

We say that γ' is obtained from γ by a *SL-move*. For example, see Figure 3.8. We note that if u is trivial, a SL-move is just adding full-twists or nothing. A SL-move is determined by a string link and a number of full-twists, that is, ‘SL’ stands for String Link.

Proposition 3.1. Let γ be an n -component bottom tangle and γ' a bottom tangle obtained from γ by a SL-move. If the Milnor numbers of γ and γ' for sequences with length $\leq k$ vanish, then for any sequence I ,

$$\mu_{\gamma'}(I) \equiv \mu_{\gamma}(I) \pmod{\delta_{\gamma}^k(I)},$$

where $\delta_{\gamma}^k(I)$ is the greatest common divisor of $\mu_{\gamma}(J)$'s for a proper subsequence J of I which is obtained by removing at least $k + 1$ indices.




FIGURE 3.8. An example of a SL-move

Proof. The proof is by induction on the length $|I| = q$. If $q \leq k$, the proposition clearly holds. We note that, by the induction hypothesis, $\delta_\gamma^k(I) = \delta_{\gamma'}^k(I)$. Denote respectively by α_i, λ_i (resp. α'_i, λ'_i) the i th meridian and i th longitude of γ (resp. γ') for $1 \leq i \leq n$. Let E_X (resp. E_Y) be the Magnus expansion in non-commutative variables $X_1, \dots, X_n$ (resp. $Y_1, \dots, Y_n$) obtained by replacing α_i by $1 + X_i$ (resp. α'_i by $1 + Y_i$) for $1 \leq i \leq n$. Fix j , by the assumption, the Milnor numbers for γ and γ' of length $\leq k$ vanish, so $E_X(\lambda_j)$ and $E_Y(\lambda'_j)$ can be written respectively in the form

$$E_X(\lambda_j) = 1 + F_j(X) \text{ and } E_Y(\lambda'_j) = 1 + F'_j(Y),$$

where $F_j(X) (= F_j(X_1, \dots, X_n))$ and $F'_j(Y) (= F'_j(Y_1, \dots, Y_n))$ are terms of degree $\geq k$.

Here we define a set of polynomials;

$$D_j^k = \left\{ \sum \nu(i_1 \dots i_m) Y_{i_1} \dots Y_{i_m} \mid \begin{array}{ll} \nu(i_1 \dots i_m) \equiv 0 \pmod{\delta_\gamma^k(i_1 \dots i_m j)}, & m < q \\ \nu(i_1 \dots i_m) \in \mathbb{Z}, & m \geq q \end{array} \right\}.$$

Then it is enough to show

$$F'_j(Y) - F_j(Y) \in D_j^k.$$

The following claims are shown by similar to the assertions (16) and (18) in [8].

Claim 1. D_j^k is a two-sided ideal of the formal power series ring in non-commutative variables $Y_1, \dots, Y_n$ with integer coefficients.

Claim 2. If at least k variables are inserted anywhere in a term $\mu(i_1 i_2 \dots i_m j) Y_{i_1 i_2 \dots i_m}$, then the resulting term belongs to D_j^k .

Let u_i be the i th longitude of a string link which gives the SL-move, see Figure 3.9. In the proof of [10, Lemma 2.6], it is shown that the degree of each term in $E_Y(u_i^{\pm 1}) - 1$ is at least $k + 1$. Set $E_Y(u_i) = 1 + G_i(Y)$ and $E_Y(u_i^{-1}) = 1 + \overline{G}_i(Y)$, where $G_i(Y)$ and $\overline{G}_i(Y)$ mean the terms of degree $\geq k + 1$. Since $\alpha_i = u_i^{-1} \alpha'_i u_i$ (where α_i is assumed to be an element of $\pi_1([0, 1]^3 \setminus \gamma')$), we have

$$\begin{aligned} E_Y(\alpha_i) &= E_Y(u_i^{-1} \alpha'_i u_i) \\ &= (1 + \overline{G}_i(Y))(1 + Y_i)(1 + G_i(Y)) \\ &= (1 + \overline{G}_i(Y))(1 + G_i(Y)) + (1 + \overline{G}_i(Y))Y_i(1 + G_i(Y)) \\ &= 1 + Y_i + Y_i G_i(Y) + \overline{G}_i(Y)Y_i + \overline{G}_i(Y)Y_i G_i(Y). \end{aligned}$$

Hence $E_Y(\lambda_j)$ is obtained from $E_X(\lambda_j)$ by substituting X_i for

$$Y_i + Y_i G_i(Y) + \overline{G}_i(Y)Y_i + \overline{G}_i(Y)Y_i G_i(Y).$$

Set $E_Y(\lambda_j) = 1 + H_j(Y)$, where $H_j(Y)$ is the terms of degree $\geq k$. Note that terms of degree $\leq 2k$ of $H_j(Y) - F_j(Y)$ vanish, and that any term of $H_j(Y) - F_j(Y)$ of




FIGURE 3.9.

degree $\geq 2k + 1$ is obtained from $F_j(Y)$ by inserting at least $k + 1$ variables. By Claim 2,

$$H_j(Y) - F_j(Y) \in D_j^k.$$

Since $\lambda'_j = u_j \lambda_j u_j^{-1}$ (where λ_j is assumed to be an element of $\pi_1([0, 1]^3 \setminus \gamma')$),

$$\begin{aligned} E_Y(\lambda'_j) &= E_Y(u_j \lambda_j u_j^{-1}) \\ &= (1 + G_j(Y))(1 + H_j(Y))(1 + \overline{G_j}(Y)) \\ &= 1 + H_j(Y) + H_j(Y)\overline{G_j}(Y) + G_j(Y)H_j(Y) + G_j(Y)H_j(Y)\overline{G_j}(Y). \end{aligned}$$

It follows from Claims 1 and 2 that we have

$$F'_j(Y) - H_j(Y) = H_j(Y)\overline{G_j}(Y) + G_j(Y)H_j(Y) + G_j(Y)H_j(Y)\overline{G_j}(Y) \in D_j^k.$$

Since $H_j(Y) - F_j(Y) \in D_j^k$, by Claim 1, we have

$$F'_j(Y) - F_j(Y) \in D_j^k.$$

This completes the proof. $\square$

Proof of Theorem 1.1. By Lemma 2.1, any two disk/band surfaces F_c and F'_c of an n -component clover link c are transformed into each other by the moves (a) and (b) illustrated in Figure 2.4. So two bottom tangles $\gamma(F_c)$ and $\gamma(F'_c)$ are transformed into each other by a SL-move. Since the both closures $L(\gamma(F_c))$ and $L(\gamma(F'_c))$ are ambient isotopic to l_c , by the hypothesis of Theorem 1.1,

$$0 = \overline{\mu}_{l_c}(J) = \mu_{\gamma(F_c)}(J) = \mu_{\gamma(F'_c)}(J)$$

for any sequence J with $|J| \leq k$. Hence by Proposition 3.1,

$$\mu_{\gamma(F'_c)}(I) \equiv \mu_{\gamma(F_c)}(I) \pmod{\delta_{\gamma(F_c)}^k(I)}$$

for any sequence I . This completes the proof. $\square$

4. PROOF OF THEOREMS 1.3 AND 1.6

In this section we will give proofs of Theorems 1.3, 1.6 and Corollary 1.4.

Proposition 4.1. *Let γ be an n -component bottom tangle and γ' a bottom tangle obtained from γ by a SL-move which is given by a string link u . If the Milnor numbers of γ and γ' for non-repeated sequences with length $\leq k$ vanish, then we have the following:*

$$\begin{aligned}
& \sum_{S \in \mathcal{S}_j^{2k+1}} (\mu_{\gamma'}(Sj) - \mu_{\gamma}(Sj)) Y_S \\
= & \sum_{\substack{|J|=|I|=k \\ JI \in \mathcal{S}_j^{2k+1}}} \mu_{\gamma}(Jj) \mu_{\gamma}(Il) \sum_{i_s \in \{J\}} \mu_u(li_s) Y_{J < s} (Y_{i_s Il} - Y_{i_s IJ} - Y_{Il i_s} + Y_{I I i_s}) Y_{J < s} \\
& + \sum_{\substack{|J|=|I|=k \\ JI \in \mathcal{S}_j^{2k+1}}} \mu_{\gamma}(Jj) \mu_{\gamma}(Il) \mu_u(lj) (Y_{IlJ} - Y_{IJ} - Y_{JIl} + Y_{JII}).
\end{aligned}$$

Proof. We compare with the Magnus expansions of the j th longitudes of γ and γ' . Denote respectively by α_i, λ_i (resp. α'_i, λ'_i) the i th meridian and i th longitude of γ (resp. γ') for $1 \leq i \leq n$. Let E_X (resp. E_Y) be the Magnus expansion in non-commutative variables $X_1, \dots, X_n$ (resp. $Y_1, \dots, Y_n$) obtained by replacing α_i by $1 + X_i$ (resp. α'_i by $1 + Y_i$) for $1 \leq i \leq n$. By the assumption, the Minor numbers for γ and γ' of degree k coincide, hence denote by

$$E_X(\lambda_j) = 1 + \sum_{I \in \mathcal{S}_j^k} \mu_{\gamma}(Ij) X_I + r_j(X) + \mathcal{O}_X(2)$$

and

$$E_Y(\lambda'_j) = 1 + \sum_{I \in \mathcal{S}_j^k} \mu_{\gamma}(Ij) Y_I + r'_j(Y) + \mathcal{O}_Y(2),$$

where $r_j(X)$ and $r'_j(Y)$ mean the terms of degree $\geq k+1$ and $\mathcal{O}_X(2)$ (resp. $\mathcal{O}_Y(2)$) denotes the terms which contain X_i (resp. Y_i) at least 2 times for some $i (= 1, 2, \dots, n)$. Let $f_j(X) = \sum_{I \in \mathcal{S}_j^k} \mu_{\gamma}(Ij) X_I$ and $f_j(Y) = \sum_{I \in \mathcal{S}_j^k} \mu_{\gamma}(Ij) Y_I$.

Let u_i be the i th longitude of u and let $\beta_i = [\lambda'_i, \alpha'_i] = \lambda'^{-1}_i \alpha'^{-1}_i \lambda'_i \alpha'_i$, see Figure 3.9. Let E_Z be the Magnus expansion in non-commutative variables $Z_1, \dots, Z_n$ obtained by replacing β_i by $1 + Z_i$ for $1 \leq i \leq n$. Then we have

$$E_Z(u_i) = 1 + \sum_{l \neq i} \mu_u(li) Z_l + \mathcal{O}(2).$$

First, we observe $E_Y(\beta_l)$ and $E_Y(\alpha_i)$, where α_i is assumed to be an element of $\pi_1([0, 1]^3 \setminus \gamma')$. Since $E_Y(\lambda'_l) E_Y(\lambda'^{-1}_l) = 1$, set $E_Y(\lambda'^{-1}_l) = 1 - f_l(Y) + \overline{r'_l}(Y) + \mathcal{O}_Y(2)$, where $\overline{r'_l}(Y)$ is the terms of degree $\geq k+1$. Observe that

$$\begin{aligned}
& E_Y(\beta_l) \\
= & E_Y(\lambda'^{-1}_l \alpha'^{-1}_l \lambda'_l \alpha'_l) \\
= & (1 - f_l(Y) + \overline{r'_l}(Y) + \mathcal{O}_Y(2))(1 - Y_l + \mathcal{O}_Y(2))(1 + f_l(Y) + r'_l(Y) + \mathcal{O}_Y(2))(1 + Y_l) \\
= & 1 + f_l(Y) Y_l - Y_l f_l(Y) + \mathcal{O}(k+2) + \mathcal{O}_Y(2) \\
= & 1 + \sum_{I \in \mathcal{S}_l^k} \mu_{\gamma}(Il) Y_I Y_l - Y_l \sum_{I \in \mathcal{S}_l^k} \mu_{\gamma}(Il) Y_I + \mathcal{O}(k+2) + \mathcal{O}_Y(2) \\
= & 1 + \sum_{I \in \mathcal{S}_l^k} \mu_{\gamma}(Il) (Y_{Il} - Y_{II}) + \mathcal{O}(k+2) + \mathcal{O}_Y(2).
\end{aligned}$$

This implies that $E_Y(u_i)$ is obtained from $E_Z(u_i)$ by substituting Z_l for

$$\sum_{I \in \mathcal{S}_l^k} \mu_{\gamma}(Il) (Y_{Il} - Y_{II}) + \mathcal{O}(k+2) + \mathcal{O}_Y(2).$$

So we have

$$E_Y(u_i) = 1 + \sum_{l \neq i} \mu_u(li) \sum_{I \in \mathcal{S}_l^k} \mu_{\gamma}(Il) (Y_{Il} - Y_{II}) + \mathcal{O}(k+2) + \mathcal{O}_Y(2).$$

Let $g_i(Y) = \sum_{l \neq i} \mu_u(li) \sum_{I \in \mathcal{S}_l^k} \mu_\gamma(I) (Y_{Il} - Y_{lI})$. Then we have

$$E_Y(u_i^{-1}) = 1 - g_i(Y) + \mathcal{O}(k+2) + \mathcal{O}_Y(2).$$

Since $\alpha_i = u_i^{-1} \alpha'_i u_i$, we have

$$\begin{aligned} E_Y(\alpha_i) &= E_Y(u_i^{-1} \alpha'_i u_i) \\ &= (1 - g_i(Y) + \mathcal{O}(k+2) + \mathcal{O}_Y(2))(1 + Y_i)(1 + g_i(Y) + \mathcal{O}(k+2) + \mathcal{O}_Y(2)) \\ &= 1 + Y_i + Y_i g_i(Y) - g_i(Y) Y_i + \mathcal{O}(k+3) + \mathcal{O}_Y(2) \\ &= 1 + Y_i + \sum_{l \neq i} \mu_u(li) \sum_{I \in \mathcal{S}_l^k} \mu_\gamma(I) (Y_{iIl} - Y_{iIl} - Y_{lIi} + Y_{lIi}) + \mathcal{O}(k+3) + \mathcal{O}_Y(2). \end{aligned}$$

Now we consider the difference $d_j(Y) = E_Y(\lambda_j) - (1 + f_j(Y) + r_j(Y) + \mathcal{O}_Y(2))$. Since $E_Y(\lambda_j)$ is obtained from $E_X(\lambda_j) (= 1 + f_j(X) + r_j(X) + \mathcal{O}_X(2))$ by substituting X_i for

$$Y_i + \sum_{l \neq i} \mu_u(li) \sum_{I \in \mathcal{S}_l^k} \mu_\gamma(I) (Y_{iIl} - Y_{iIl} - Y_{lIi} + Y_{lIi}) + \mathcal{O}(k+3) + \mathcal{O}_Y(2),$$

all terms of degree $\leq 2k$ of $d_j(Y) - \mathcal{O}_Y(2)$ vanish. The terms of degree $2k+1$ in $d_j(Y) - \mathcal{O}_Y(2)$ is obtained from $f_j(Y)$ by substituting Y_i for

$$\sum_{l \neq i} \mu_u(li) \sum_{I \in \mathcal{S}_l^k} \mu_\gamma(I) (Y_{iIl} - Y_{iIl} - Y_{lIi} + Y_{lIi})$$

for some $i \in \{1, 2, \dots, n\}$. It follows that

$$\begin{aligned} &d_j(Y) - (\mathcal{O}_Y(2) + \mathcal{O}(2k+2) + \mathcal{O}_{Y_j}) \\ &= \sum_{J \in \mathcal{S}_j^k} \mu_\gamma(J) \sum_{i_s \in \{J\}} Y_{J_{<s}} \left(\sum_{l \neq i_s} \mu_u(li_s) \sum_{JIl \in \mathcal{S}_j^{2k+1}} \mu_\gamma(I) (Y_{i_s Il} - Y_{i_s lI} - Y_{lI i_s} + Y_{lI i_s}) \right) Y_{J_{s <}} \\ &= \sum_{\substack{|J|=|I|=k \\ JIl \in \mathcal{S}_j^{2k+1}}} \mu_\gamma(J) \mu_\gamma(I) \sum_{i_s \in \{J\}} \mu_u(li_s) Y_{J_{<s}} (Y_{i_s Il} - Y_{i_s lI} - Y_{lI i_s} + Y_{lI i_s}) Y_{J_{s <}}, \end{aligned}$$

where $\mathcal{O}_{Y_j}$ means the terms which contain Y_j at least one time.

Finally, we observe the difference $E_Y(\lambda'_j) - (1 + f_j(Y) + r_j(Y) + \mathcal{O}_Y(2))$. Since $\lambda'_j = u_j \lambda_j u_j^{-1}$ (where λ_j is assumed to be an element of $\pi_1([0, 1]^3 \setminus \gamma')$), we have

$$\begin{aligned} E_Y(\lambda'_j) &= E_Y(u_j \lambda_j u_j^{-1}) \\ &= 1 + (1 + g_j(Y))(f_j(Y) + r_j(Y) + d_j(Y))(1 - g_j(Y)) + \mathcal{O}(2k+2) + \mathcal{O}_Y(2) \\ &= 1 + f_j(Y) + r_j(Y) + d_j(Y) + g_j(Y) f_j(Y) - f_j(Y) g_j(Y) + \mathcal{O}(2k+2) + \mathcal{O}_Y(2). \end{aligned}$$

So we have

$$\begin{aligned} &E_Y(\lambda'_j) - (1 + f_j(Y) + r_j(Y) + \mathcal{O}_Y(2)) \\ &= d_j(Y) + \sum_{\substack{|J|=|I|=k \\ JIl \in \mathcal{S}_j^{2k+1}}} \mu_\gamma(J) \mu_\gamma(I) \mu_u(lj) (Y_{IlJ} - Y_{lIJ} - Y_{JIl} + Y_{JI l}) \\ &\quad + \mathcal{O}(2k+2) + \mathcal{O}_Y(2) \\ &= \sum_{\substack{|J|=|I|=k \\ JIl \in \mathcal{S}_j^{2k+1}}} \mu_\gamma(J) \mu_\gamma(I) \sum_{i_s \in \{J\}} \mu_u(li_s) Y_{J_{<s}} (Y_{i_s Il} - Y_{i_s lI} - Y_{lI i_s} + Y_{lI i_s}) Y_{J_{s <}} \\ &\quad + \sum_{\substack{|J|=|I|=k \\ JIl \in \mathcal{S}_j^{2k+1}}} \mu_\gamma(J) \mu_\gamma(I) \mu_u(lj) (Y_{IlJ} - Y_{lIJ} - Y_{JIl} + Y_{JI l}) \\ &\quad + \mathcal{O}(2k+2) + \mathcal{O}_Y(2) + \mathcal{O}_{Y_j}. \end{aligned}$$

This completes the proof. $\square$

Proof of Theorem 1.3. By Lemma 2.1, any two disk/band surfaces F_c and F'_c of an n -component clover link c are transformed into each other by the moves (a) and (b) in Figure 2.4. So two bottom tangles $\gamma(F_c)$ and $\gamma(F'_c)$ are transformed into each other by a SL-move. Since the both closures $L(\gamma(F_c))$ and $L(\gamma(F'_c))$ are ambient isotopic to l_c and the hypothesis of Theorem 1.3,

$$0 = \overline{\mu}_{l_c}(J) = \mu_{\gamma(F_c)}(J) = \mu_{\gamma(F'_c)}(J)$$

for any sequence J with $|J| \leq k$. Since $\mu_u(pq)$ is the ‘linking number’ of the p th component and the q th component of u , $\mu_u(pq) = \mu_u(qp)$ and the set

$$\{\mu_u(pq) \mid u : \text{a string link}\} = \mathbb{Z}$$

for any p and q . This and Proposition 4.1 give us the Theorem 1.3. $\square$

In order to prove Corollary 1.4 and Theorem 1.6, we need the following lemma given in [10].

Lemma 4.2. [10, Lemma 4.1] *Two n -clover links c and c' are edge-homotopic if and only if there exist disk/band surfaces F_c and $F_{c'}$ of c and c' respectively such that the two bottom tangles $\gamma(F_c)$ and $\gamma(F_{c'})$ are link-homotopic.*

Proof of Corollary 1.4. Let c and c' be n -clover links. We assume that they are edge-homotopic. By Lemma 4.2, there exist disk/band surfaces F_c and $F_{c'}$ of c and c' respectively such that $\gamma(F_c)$ and $\gamma(F_{c'})$ are link-homotopic. This implies that $\mu_{\gamma(F_c)}(I) = \mu_{\gamma(F_{c'})}(I)$ for any non-repeated sequence I [7], [2]. On the other hand, by Theorem 1.3, the set $H_c(2k+2, j)$ (resp. $H_{c'}(2k+2, j)$) is obtained from the Milnor numbers of $\gamma(F_c)$ (resp. $\gamma(F_{c'})$) for F_c (resp. $F_{c'}$). Hence we have $H_c(2k+2, j) = H_{c'}(2k+2, j)$. $\square$

Proof of Theorem 1.6. Suppose that two 4-clover links c and c' are edge-homotopic. By Corollary 1.4, $H_c(4, 4) = H_{c'}(4, 4) (\neq \emptyset)$. By Lemma 4.2 there exist disk/band surfaces F_c and $F_{c'}$ such that $\gamma(F_c)$ and $\gamma(F_{c'})$ are link-homotopic. This implies that the Milnor numbers of $\gamma(F_c)$ and $\gamma(F_{c'})$ are equal for any non-repeated sequence [7], [2]. Since the Milnor numbers of length ≤ 3 are always well-defined for clover links (see Remark 2.3), we have $\mu_c(I) = \mu_{c'}(I)$ for any non-repeated sequence I with $|I| \leq 3$.

Conversely if $H_c(4, 4) \cap H_{c'}(4, 4) \neq \emptyset$, then there exist disk/band surfaces F_c and $F_{c'}$ of c and c' respectively such that

$$\sum_{S \in \mathcal{S}_4^3} \mu_{\gamma(F_c)}(S4) X_S = \sum_{S \in \mathcal{S}_4^3} \mu_{\gamma(F_{c'})}(S4) X_S.$$

In particular,

$$\mu_{\gamma(F_c)}(1234) = \mu_{\gamma(F_{c'})}(1234) \text{ and } \mu_{\gamma(F_c)}(2134) = \mu_{\gamma(F_{c'})}(2134).$$

According to the link-homotopy classification theorem for string links by N. Habegger and X. S. Lin [2], for two 4-component string links (bottom tangles) that have common values of the Milnor numbers for non-repeated sequences with length ≤ 3 , they are link-homotopic if and only if their Milnor numbers for sequences 1234 and 2134 coincide, see also [11, Theorem 4.3]. This together with the hypothesis implies that $\gamma(F_c)$ and $\gamma(F_{c'})$ are link-homotopic. Therefore c and c' are edge-homotopic by Lemma 4.2. This completes the proof. $\square$

REFERENCES

- [1] K. T. Chen, *Commutator calculus and link invariants*, Proc. Amer. Math. Soc. 3 (1952), 44–55.
- [2] N. Habegger; X. S. Lin, *The classification of links up to link-homotopy*, J. Amer. Math. Soc. 3 (1990), 389–419.
- [3] K. Habiro, *Bottom tangles and universal invariants*, Algebr. Geom. Topol. 6 (2006), 1113–1214.
- [4] L. Kauffman; J. Simon; K. Wolcott; P. Zhao, *Invariants of theta-curves and other graphs in 3-space*, Topology Appl. 49 (1993), 193–216.
- [5] J. LeDimet, *Cobordisme d'enlacements de disques* (French), Mém. Soc. Math. France (N.S.) No. 32 (1988).
- [6] J. P. Levine, *The $\bar{\mu}$ -invariants of based links*, Differential topology (Siegen, 1987), 87–103, Lecture Notes in Math., 1350, Springer, Berlin, (1988).
- [7] J. Milnor, *Link groups*, Ann. of Math. (2) 59 (1954), 177–195.
- [8] J. Milnor, *Isotopy of links*, Algebraic geometry and topology, A symposium in honor of S. Lefschetz, pp. 280–306. Princeton University Press, Princeton, N. J. (1957).
- [9] K. Taniyama, *Cobordism, homotopy and homology of graphs in $\mathbf{R}^3$* , Topology 33 (1994), 509–523.
- [10] K. Wada, *Milnor invariants and edge-homotopy classification of clover links*, preprint (2015).
- [11] A. Yasuhara, *Self delta-equivalence for links whose Milnor's isotopy invariants vanish*, Trans. Amer. Math. Soc. 361 (2009), 4721–4749.

DEPARTMENT OF MATHEMATICS, SCHOOL OF EDUCATION, WASEDA UNIVERSITY, NISHI-WASEDA 1-6-1, SHINJUKU-KU, TOKYO, 169-8050, JAPAN

DEPARTMENT OF MATHEMATICS, TOKYO GAUGEI UNIVERSITY, 4-1-1 NUKUIKITA-MACHI, KOGANEI-SHI, TOKYO, 184-8501, JAPAN

E-mail address: k.wada@akane.waseda.jp, yasuhara@u-gakugei.ac.jp